

L5(6.1) - Solving Quadratic Equations

Recall:

To solve an equation, find value(s) that satisfy the equation (i.e., make it true).

This value is called the solution or root of the equation.

Ex.1 Solve $x^2 - 12x + 32 = 0$

Apr 19-7:41 PM

We have most often solved for the zeroes of the quadratic equation, but we can solve for any value.


Ex.2 Solve $y = 2x^2 + 5x - 12$ for

(a) $y = 0$

(b) $y = -12$

Nov 22-7:19 PM

Ex.2 Solve $y = 2x^2 + 5x - 12$ for (a) $y = 0$
(b) $y = -12$



Nov 22-7:19 PM


To solve using factored form:

- 1) Expand all terms
- 2) Move all terms to one side of the equal sign so that the equation equals zero
- 3) Factor your expression (if possible)
- 4) Set each factor equal to zero and solve

Ex.3 Solve: $x^2 - 10 = -x(2x + 13)$


Mar 29-11:52 AM

$$y = x^2 - 10$$



Nov 22-7:51 PM

$$y = -x(2x + 13)$$




Nov 22-7:51 PM

$$x^2 - 10 = -x(2x + 13)$$

which became

$$3x^2 + 13x - 10 = 0$$

$$(x + 5)(3x - 2) = 0$$



Nov 22-7:51 PM

Assigned Work:

p.320 # 4ac, 6ace, 7ace, 9ace, 11, 14